ADDENDUM TO TABLE 9, PAGE 30

Please note the Strata-bound Type includes the Upper Granite Breccia Member and the Brooks Member as well as the Whenan Member and the Black Hematite Member. The Transgressive Type includes the Lower Silicified Member, the Upper Granite Breccia and the Brooks Member.

Table 9

MINERALIZED FORMATIONS AT OLYMPIC DAM[1]

Geologic Age — Middle Proterozoic
Mineralized Sequence
Strata-bound Type
Transgressive Type

Greenfield Formation	
Volcanic Member (250 m thick)	Volcanic breccias and conglomerates
Hematite Breccia Member (500 m thick)	Hematite-rich breccias and siltstones
Lower Silicified Member (275 m)	Hematite and silicifed breccias, *chalcocite* and *bornite minerals*
Olympic Dam Formation	
Upper Granite Breccia Member (100 m thick)	Hematite-rich granite breccia, *chalcocite* and *bornite minerals*
Brooks Member (300 m thick)	Mixed rock breccias, brown and black hematite matrix, *chalcopyrite, bornite and chalcocite minerals*
Whenan Member (350 m thick)	Mixed rock breccias, black and red hematite matrix, bedded units throughout, *bornite* and *chalcopyrite minerals*
Black Hematite Member (150 m thick)	Mixed rock breccias, black hematite matrix, *chalcopyrite mineral*
Lower Granite Breccia Member (800 m thick)	Granite and mixed rock breccias, chlorite alteration

[1]Adapted from D.E. Roberts and G.R.T. Hudson, RMS, Economic Geology, August 1983.

© NUEXCO 1983